arXiv:1702.06420v1 [math.NT] 20 Feb 2017

A closed formula for the generating function of p-Bernoulli numbers

Levent Kargın[1] and Mourad Rahmani[2]
[1]Akseki Vocational School, Alanya Alaaddin Keykubat University, Antalya TR-07630, Turkey
[2]USTHB, Faculty of Mathematics, P. O. Box32, El Alia, Bab Ezzouar, 16111, Algiers, Algeria
leventkargin48@gmail.com and mrahmani@usthb.dz

Abstract

In this paper, using geometric polynomials, we obtain a generating function of p-Bernoulli numbers. As a consequences this generating function, we derive closed formulas for the finite summation of Bernoulli and harmonic numbers involving Stirling numbers of the second kind.

2000 Mathematics Subject Classification: 11B68, 11B83

Key words: p-Bernoulli number, geometric polynomial, harmonic number.

1 Introduction

Rahmani [9] introduced p-Bernoulli numbers by constructing an infinite matrix as follows:

The first row of the matrix $B_{0,p}=1$ and each entry $B_{n,p}$ is given recursively by

$$B_{n+1,p}=pB_{n,p}-\frac{(p+1)^2}{p+2}B_{n,p+1}.$$

The first column of the matrix $B_{n,0}=B_n$. Here, B_n is the nth Bernoulli number.

For every integer $p\geq -1$, these numbers have an explicit formula

$$B_{n,p}=\sum_{k=0}^{n}(-1)^k\left\{{n \atop k}\right\}\binom{k+p+1}{k}^{-1}k!, \tag{1}$$

and are closely related to Bernoulli numbers by the following formula

$$B_{n,p}=\frac{p+1}{p!}\sum_{j=0}^{p}(-1)^j\left[{p \atop j}\right]B_{n+j}, \tag{2}$$

where $\begin{bmatrix} n \\ k \end{bmatrix}$ is the Stirling number of the first kind [6]. The first few generating functions for $B_{n,p}$ $(p = 1, 2)$ are

$$\sum_{n=0}^{\infty} B_{n,1} \frac{t^n}{n!} = \frac{2\left[(t-1)\, e^t + 1\right]}{\left(e^t - 1\right)^2},$$
$$\sum_{n=0}^{\infty} B_{n,2} \frac{t^n}{n!} = \frac{3\left[(2t-3)\, e^{2t} + 4e^t - 1\right]}{2\left(e^t - 1\right)^3}.$$

The main purpose of this study is to give a close form of the above results as

$$\sum_{n=0}^{\infty} B_{n,p} \frac{t^n}{n!} = \frac{(p+1)\,(t - H_p)\, e^{pt}}{(e^t - 1)^{p+1}} + (p+1) \sum_{k=1}^{p} \binom{p}{k} \frac{H_k}{(e^t - 1)^{k+1}}, \tag{3}$$

where H_n is the harmonic numbers, defined by [6, p. 258]

$$H_n = \sum_{j=1}^{n} \frac{1}{j}.$$

As a consequences of (3), we have closed formulas for the finite summation of Bernoulli and harmonic numbers.

For the proof of (3), we use some properties of geometric polynomials. The geometric polynomials are defined by means of the following generating function [10]

$$\frac{1}{1 - x\left(e^t - 1\right)} = \sum_{n=0}^{\infty} w_n(x) \frac{t^n}{n!}, \tag{4}$$

and have the explicit formula

$$w_n(x) = \sum_{k=0}^{n} \left\{ {n \atop k} \right\} k! x^k, \tag{5}$$

where $\left\{ {n \atop k} \right\}$ is the Stirling number of the second kind [6]. The Stirling numbers of the second kind are defined by means of the following generating function

$$\sum_{n=0}^{\infty} \left\{ {n \atop k} \right\} \frac{t^n}{n!} = \frac{\left(e^t - 1\right)^k}{k!}. \tag{6}$$

Some other properties of geometric polynomials can be found in [1, 2, 3, 4, 5, 7].

2 A new generating function for p-Bernoulli numbers

In this section, the main theorem and its applications are given.

Now, we give the main theorem of this paper.

Theorem 1 *For $p \geq 0$, the following generating function holds true:*

$$\sum_{n=0}^{\infty} B_{n,p} \frac{t^n}{n!} = \frac{(p+1)\left(t - H_p\right) e^{pt}}{\left(e^t - 1\right)^{p+1}} + (p+1) \sum_{k=1}^{p} \binom{p}{k} \frac{H_k}{\left(e^t - 1\right)^{k+1}}. \tag{7}$$

For the proof of main theorem, we first need the following proposition.

Proposition 2 *For $n > p \geq 0$, we have*

$$\frac{(-1)^p}{(p+1)!} B_{n,p} = \underbrace{\int_{-1}^{0} \int_{0}^{x_p} \dots \int_{0}^{x_3} \int_{0}^{x_2}}_{p+1 \ times} w_n\left(x_1\right) dx_1 dx_2 \dots dx_{p-1} dx_p. \tag{8}$$

Proof. If we integrate both sides of (5) with respect to x_1 from 0 to x_2, we have

$$\int_{0}^{x_2} w_n\left(x_1\right) dx_1 = \sum_{k=0}^{n} \left\{ n \atop k \right\} k! \frac{x_2^{k+1}}{k+1}.$$

Integrating both sides of the above equation with respect to x_2 from 0 to x_3, we obtain

$$\int_{0}^{x_3} \int_{0}^{x_2} w_n\left(x_1\right) dx_1 dx_2 = \sum_{k=0}^{n} \left\{ n \atop k \right\} k! \frac{x_3^{k+2}}{(k+1)(k+2)}.$$

Applying the same procedure for p times yields

$$\int_{0}^{x_p} \dots \int_{0}^{x_3} \int_{0}^{x_2} w_n\left(x_1\right) dx_1 dx_2 \dots dx_{p-1} = \sum_{k=0}^{n} \left\{ n \atop k \right\} k! \frac{x_p^{k+p}}{(k+1) \cdots (k+p)}.$$

Finally, integrating both sides of the above equation with respect to x_p from -1 to 0 and using (1) gives the desired equation. ■

We note that taking $p = 0$ in (8) gives [8, Theorem 1.2].

Now, we are ready to give the proof of the main theorem.

Proof of Theorem 1. Multiplying both sides of (8) with $\frac{t^n}{n!}$ and summing over n from 0 to infinitive, we have

$$\begin{aligned} \frac{(-1)^p}{(p+1)!} \sum_{n=0}^{\infty} B_{n,p} \frac{t^n}{n!} &= \int_{-1}^{0} \int_{0}^{x_p} \dots \int_{0}^{x_3} \left[\int_{0}^{x_2} \left(\sum_{n=0}^{\infty} w_n\left(x_1\right) \frac{t^n}{n!} \right) dx_1 \right] dx_2 \dots dx_{p-1} dx_p \\ &= \int_{-1}^{0} \int_{0}^{x_p} \dots \int_{0}^{x_3} \left[\int_{0}^{x_2} \frac{1}{1 - x_1\left(e^t - 1\right)} dx_1 \right] dx_2 \dots dx_{p-1} dx_p. \end{aligned}$$

If we evaluate the first integral, we obtain

$$\int_{0}^{x_2} \frac{1}{1 - x_1\left(e^t - 1\right)} dx_1 = \frac{-1}{e^t - 1} \ln\left(1 - x_2\left(e^t - 1\right)\right).$$

For the second time, we evaluate

$$\frac{-1}{e^t-1}\int_0^{x_3}\ln\left(1-x_2\left(e^t-1\right)\right)dx_2$$
$$=\frac{1}{\left(e^t-1\right)^2}\left[\left(1-x_3\left(e^t-1\right)\right)\ln\left(1-x_3\left(e^t-1\right)\right)-\left(1-x_3\left(e^t-1\right)\right)+1\right].$$

By induction on p, let us assume that the following equation holds

$$\int_0^{x_p}\cdots\int_0^{x_3}\int_0^{x_2}w_n\left(x_1\right)dx_1dx_2\ldots dx_{p-1} \tag{9}$$
$$=\frac{(-1)^p}{\left(e^t-1\right)^p(p-1)!}\left(1-x_p\left(e^t-1\right)\right)^{p-1}\ln\left(1-x_p\left(e^t-1\right)\right)$$
$$-\frac{(-1)^p}{\left(e^t-1\right)^p(p-1)!}\left[H_{p-1}\left(1-x_p\left(e^t-1\right)\right)^{p-1}+H_{p-1}\right]$$
$$+\sum_{k=1}^{p-2}\frac{(-1)^{p-k}H_{p-1-k}x_p^k}{\left(e^t-1\right)^{p-k}(p-1-k)!k!}.$$

Now, we want to prove that (9) holds for the case $p+1$. Let us integrate both sides of (9) with respect to x_p from 0 to y. Then we have

$$\int_0^{y}\int_0^{x_p}\cdots\int_0^{x_3}\int_0^{x_2}w_n\left(x_1\right)dx_1dx_2\ldots dx_{p-1}dx_p$$
$$=\frac{(-1)^p}{\left(e^t-1\right)^p(p-1)!}\int_0^{y}\left(1-x_p\left(e^t-1\right)\right)^{p-1}\ln\left(1-x_p\left(e^t-1\right)\right)dx_p$$
$$-\frac{(-1)^pH_{p-1}}{\left(e^t-1\right)^p(p-1)!}\int_0^{y}\left(1-x_p\left(e^t-1\right)\right)^{p-1}dx_p+\frac{(-1)^pH_{p-1}}{\left(e^t-1\right)^p(p-1)!}\int_0^{y}dx_p$$
$$+\sum_{k=1}^{p-2}\frac{(-1)^{p-k}H_{p-1-k}}{\left(e^t-1\right)^{p-k}(p-1-k)!k!}\int_0^{y}x_p^kdx_p.$$

The first integral in the right hand-side equals

$$\frac{(-1)^p}{\left(e^t-1\right)^p(p-1)!}\int_0^{y}\left(1-x_p\left(e^t-1\right)\right)^{p-1}\ln\left(1-x_p\left(e^t-1\right)\right)dx_p \tag{10}$$
$$=\frac{(-1)^{p+1}}{\left(e^t-1\right)^{p+1}p!}\left[\left(1-y\left(e^t-1\right)\right)^p\ln\left(1-y\left(e^t-1\right)\right)-\frac{\left(1-y\left(e^t-1\right)\right)^p}{p}+\frac{1}{p}\right].$$

For the second integral in the right hand-side, we obtain

$$\begin{aligned}&\frac{(-1)^p H_{p-1}}{(e^t-1)^p (p-1)!}\int_0^y \left(1-x_p\left(e^t-1\right)\right)^{p-1} dx_p \qquad (11)\\ &= \frac{(-1)^{p+1} H_{p-1}}{(e^t-1)^{p+1} p!}\left(1-y\left(e^t-1\right)\right)^p - \frac{(-1)^{p+1} H_{p-1}}{(e^t-1)^{p+1} p!}.\end{aligned}$$

For the third and fourth integrals, we find

$$\frac{(-1)^p H_{p-1}}{(e^t-1)^p (p-1)!}\int_0^y dx_p = \frac{(-1)^{p+1} H_{p-1}}{(e^t-1)^p (p-1)!} y \qquad (12)$$

and

$$\sum_{k=1}^{p-2} \frac{(-1)^{p-k} H_{p-1-k}}{(e^t-1)^{p-k} (p-1-k)!k!}\int_0^y x_p^k dx_p = \sum_{k=2}^{p-1} \frac{(-1)^{p+1-k} H_{p-k} y^k}{(e^t-1)^{p+1-k} (p-k)!k!}, \qquad (13)$$

respectively. Combining (10), (11), (12) and (13), we achieve that

$$\begin{aligned}&\int_0^y\int_0^{x_p}\cdots\int_0^{x_3}\int_0^{x_2} w_n\left(x_1\right) dx_1 dx_2 \ldots dx_{p-1} dx_p\\ &= \frac{(-1)^{p+1}}{(e^t-1)^{p+1} p!}\left(1-y\left(e^t-1\right)\right)^p \ln\left(1-y\left(e^t-1\right)\right)\\ &- \frac{(-1)^{p+1}}{(e^t-1)^{p+1} p!}\left[H_p\left(1-y\left(e^t-1\right)\right)^p + H_p\right]\\ &+ \sum_{k=1}^{p-1} \frac{(-1)^{p+1-k} H_{p-k} y^k}{(e^t-1)^{p+1-k} (p-k)!k!}.\end{aligned}$$

Finally, setting $y=-1$ in the above equation and using (8), we arrive at the desired equation. ■

As an application of Theorem 1, we give the following theorem.

Theorem 3 *For $n > p \geq 0$, we have*

$$\sum_{k=p+1}^{n} \binom{n}{k} \left\{ {k \atop p+1} \right\} B_{n-k,p} = \frac{p^{n-1}\left(n-pH_p\right)}{p!} + \sum_{j=1}^{p} \left\{ {n \atop p-j} \right\} \frac{H_j}{j!}. \qquad (14)$$

Proof. Multiplying both sides of (7) with $\frac{\left(e^t-1\right)^{p+1}}{(p+1)!}$ and using (6), the left hand

side of (7) becomes

$$\frac{(e^t-1)^{p+1}}{(p+1)!}\sum_{n=0}^{\infty}B_{n,p}\frac{t^n}{n!}=\sum_{n=0}^{\infty}\sum_{k=0}^{\infty}\left\{ {k \atop p+1}\right\}\frac{B_{n,p}}{k!n!}t^{n+k} \tag{15}$$
$$=\sum_{n=0}^{\infty}\left[\sum_{k=0}^{n}\binom{n}{k}\left\{ {k \atop p+1}\right\}B_{n-k,p}\right]\frac{t^n}{n!}.$$

For the right hand side of (7), we obtain

$$\frac{te^{pt}}{p!}-\frac{H_pe^{pt}}{p!}+\sum_{k=1}^{p}\frac{H_k}{k!}\frac{(e^t-1)^{p-k}}{(p-k)!} \tag{16}$$
$$=\sum_{n=1}^{\infty}\left[\frac{p^{n-1}(n-pH_p)}{p!}+\sum_{k=1}^{p}\frac{H_k}{k!}\left\{ {n \atop p-k}\right\}\right]\frac{t^n}{n!}.$$

Finally, comparing the coefficients of $\frac{t^n}{n!}$ in (15) and (16) completes the proof. ■

Using (2) in Theorem 3 gives the following corollary.

Corollary 4 *For $n>p\geq 0$,*

$$\sum_{k=p+1}^{n}\sum_{j=0}^{p}\binom{n}{k}\left\{ {k \atop p+1}\right\}\begin{bmatrix}p\\ j\end{bmatrix}(-1)^j B_{n+j-k}=\frac{p^{n-1}(n-pH_p)}{p+1}+\frac{p!}{p+1}\sum_{j=1}^{p}\left\{ {n \atop p-j}\right\}\frac{H_j}{j!}.$$

As a consequences of Corollary 4, the following sums are obtained $(p=1,2)$:

$$\sum_{k=2}^{n}\binom{n}{k}\left\{ {k \atop 2}\right\}B_{n+1-k}=\frac{-(n-1)}{2},$$
$$\sum_{k=3}^{n}\binom{n}{k}\left\{ {k \atop 3}\right\}(B_{n+2-k}-B_{n+1-k})=\frac{2^{n-1}(n-3)+2}{3}.$$

Setting $n=p+1$ in Theorem 3 and using $B_{0,p}=1$, we arrive at the following corollary.

Corollary 5 *For $p\geq 1$, we obtain a new closed formula for the finite summation of harmonic numbers*

$$\sum_{j=1}^{p}\left\{ {p+1 \atop p-j}\right\}\frac{H_j}{j!}=\frac{p!-(p+1)p^p+p^{p+1}H_p}{p!}.$$

References

[1] K. N. Boyadzhiev, A series transformation formula and related polynomials, *Int. J. Math. Math. Sci.* **23** (2005), 3849–3866.

[2] K. N. Boyadzhiev, Apostol-Bernoulli functions, derivative polynomials and Eulerian polynomials, *Adv. Appl. Discrete Math.* **1/2** (2008), 109–122.

[3] K. N. Boyadzhiev, Close encounters with the Stirling numbers of the second kind, *Math. Mag.* **85** (2012), 252–266.

[4] K. N. Boyadzhiev and A. Dil, Geometric polynomials: properties and applications to series with zeta values, *Anal. Math.*, **42/3** (2016), 203–224.

[5] A. Dil and V. Kurt, Investigating geometric and exponential polynomials with Euler-Seidel matrices, *J. Integer Seq.* **14** (2011), Article 11.4.6.

[6] R. L. Graham, D. E. Knuth and O. Patashnik, *Concrete Mathematics*, Addison-Wesley Publ. Com., New York, 1994.

[7] L. Kargın, Some formulae for products of Fubini polynomials with applications, arXiv:1701.01023.

[8] B. C. Kellner, Identities between polynomials related to Stirling and harmonic numbers, *Integers* **14** (2014), #A54.

[9] M. Rahmani, On p-Bernoulli numbers and polynomials, Journal of Number Theory 157 (2015) 350–366.

[10] S. M. Tanny, On some numbers related to the Bell numbers, *Canad. Math. Bull.* **17** (1974), 733–738.